

$D > 2$ TOPOLOGICAL STRING

Feng Yu

*Institute for Theoretical Physics, State University of New York**Stony Brook, New York 11794, U.S.A.**and**Department of Physics, University of Utah**Salt Lake City, Utah 84112, U.S.A.*

Abstract

A $D > 2$ topological string is presented by coupling the $2d$ topological gravity with the twisted version of the $N = 2$ superconformal matter with $c = 3k/(k - 2)$. The latter is shown to admit $k + 1$ chiral primary fields from the $SL(2, R)_k/U(1)$ unitary irreducible representations. The analysis of topological contact interactions along with the consistency requirement lead to recursion relations of correlation functions, that are convertible to the Virasoro constraints on the perturbed partition function. It is further expected to satisfy the nonlinear $\hat{W}_\infty^{(k-2)}$ constraints associated with the graded algebra $SL(k, 2)$, and thus the model is completely solvable at arbitrary genus of the surface.

Introduction — There have been encouraging new developments of string theory in last three years, led by the success of the matrix model approach[1] to the $2d$ quantum gravity, or more generally, strings with space-time dimension $D \leq 2$. The most distinguished feature is perhaps the nonperturbative solvability of all correlation functions upto all orders of Riemann surface genus, which is however best interpreted field-theoretically in the equivalent topological model approach[2]. Whereas these $D \leq 2$ models present rich and beautiful structures both physically and mathematically, in the context of string theory, they serve as toy models. It is of essential interest to search for also exactly solvable string models beyond the $D = 2$ “barrier”, where the topological phase is expected to emerge as a transition from the ordinary phase. We study the former in this Letter.

In $D < 2$ case, the relevant $2d$ topological field theory[2,3,4] – $2d$ topological gravity coupled to the twisted[5] $N = 2$ superconformal minimal matter with central charge $c = 3k/(k+2) < 3$ was shown[6] to be solved by recursion relations of physical correlators, which by consistency coincide with the extended conformal W_{k+2} constraints on the perturbed partition function. Counting in the Liouville modes of gravity sector one space-time dimension, the coupled model is usually interpreted as the $D = 1 + c/3$ topological string[7]. In the same sense, topological gravity coupled to suitable $c > 3$ conformal matter would lead to the desired $D > 2$ topological string.

There indeed exists a series of such $N = 2$ superconformal matter with $c = 3k/(k - 2)$ [8]. Naturally, its twisted version becomes the relevant topological matter, which we show in this Letter has (finitely) $k + 1$ nontrivial BRST-invariant local operators, corresponding precisely to as many chiral primaries[9] in the untwisted model. Then for the underlying $D > 2$ topological string, by analysing topological contact interactions with techniques of Ref.[4,6], slightly different recursion relations are obtained and converted to the Virasoro constraints. The consistency of multi-contact interactions involving higher-spin currents and the noncompactness of the matter sector suggests that the complete set of recursion relations, which determine all correlaton functions, are implied by the nonlinear W -constraints of the infinite-type[10,11], in particular the $\hat{W}_\infty^{(k-2)}$ constraints associated with the graded Lie algebra $SL(k, 2)$ [12].

$c > 3$ *Topological Matter* — Consider the $N = 2$ superconformal algebra, generated

by the stress tensor $T(z)$, the $U(1)$ current $J(z)$ and two spin-3/2 fermionic currents $G^\pm(z)$. For central charge $c > 3$, the $N = 2$ algebra can be realized via the generalized parafermions $\Psi^\pm(z)$ plus a free boson $\phi(z)$ [13,8], e.g.,

$$G^\pm = \sqrt{\frac{2c}{3}} \Psi^\pm e^{\pm i\sqrt{3/c}\phi}, \quad J = i\sqrt{\frac{c}{3}} \partial\phi; \quad (1)$$

so does the $SL(2, R)_k$ current algebra with

$$J^\pm = \sqrt{k} \Psi^\pm e^{\pm i\sqrt{2/k}\phi}, \quad J^3 = -\sqrt{\frac{k}{2}} \partial\phi, \quad (2)$$

where $c = 3k/(k-2)$ and the level $k(> 2)$ is restricted to integer in the scope of this Letter. Hence the irreducible representations of the former may be obtained from that of the latter. Notice the signatures of $J(z)$ in (1) and $J^3(z)$ in (2) are opposite. So although the $SL(2, R)$ module is not positive-definite because of its noncompact nature, the $N = 2$ module becomes unitary as required.

Recall there are two discrete $D^\pm$ and one continuous C series of irreducible $SL(2, R)$ representations. Indicate the lowest (highest) eigenstate of J_0^3 mode in D^+ (D^-) as $|l\rangle$ with l being positive (negative) integers or half-integers. From Ref.[8], the unitarity region for the coset $SL(2, R)_k/U(1)$ such that the $N = 2$ algebra by adding back a free boson with different signature is

$$k \geq 2|l| \geq 0, \quad k > 2; \quad (3)$$

and in the C series, simply $k > 2$. Let the primary states of the $N = 2$ representations, e.g., in the $D^\pm$ series, labeled by $|\phi_{l,n}\rangle$ ($n = 0, 1, \dots$). Note $|\phi_{l,n}\rangle$ are annihilated by positive fermion modes $G_{n+1/2}^\pm$, and the dimensions from the zero mode of $T(z)$ are

$$h = (n + \frac{1}{2})|q| - \frac{n(n+1)}{k-2} \quad (4)$$

with the $U(1)$ charges $q = -2(l \pm n)/(k-2)$ [8]. The solution of the (anti-)chiral primary states with $h = \pm q/2$ (or $G_{-1/2}^\pm |\phi_{l,n}\rangle = 0$) turns out to be $n = 0$. Thus, there are finitely many either chiral or anti-chiral primaries in D^- or D^+ series bounded by (3). Meanwhile there is no such state in the continuous series.

Upon twisting the $N = 2$ algebra (in the NS sector): $T(z) \rightarrow T(z) + (1/2)\partial J(z)$, $G_{-1/2}^+$ becomes a BRST charge Q with spin 1 while $G^-(z)$ a spin-2 current $G(z)$, such

that the stress tensor $T(z) = -\{Q, G(z)\}$ is a BRST commutator, giving the topological nature of the twisted model. The local physical operator V_α that are nontrivially BRST invariant, are precisely the chiral primary fields in the untwisted model therefore have ghost charges

$$q_\alpha = \frac{\alpha}{k-2}, \quad \alpha = 0, 1, \dots, k. \quad (5)$$

It is easy to extend the field-theoretical realization of $c < 3$ topological minimal matter with a pair of free bosons $(\varphi, \bar{\varphi})$ and free fermions $(\lambda, \bar{\lambda})$ [6] to the current $c > 3$ case, in which the (zero-form) primary operators are of the form

$$V_\alpha = e^{-\frac{\alpha}{\sqrt{k-2}}\varphi}. \quad (6)$$

The correlators of (6) and their one-form partners give rise to physical amplitudes of the underlying topological matter.

D > 2 Topological String — In turn, consider the conformal invariant description of $2d$ topological gravity[4]. Its field contents consist of the Liouville pair (ϕ, π) , their supersymmetric partner (ψ, χ) and the ghost pairs (b, c) , $(\bar{b}, \bar{c})$, (β, γ) , $(\bar{\beta}, \bar{\gamma})$ from gauge-fixing. The basic BRST-invariant (zero-form) physical operators are of the form

$$\sigma_n = \gamma_0^n = \frac{1}{2^n}(\partial\gamma + \gamma\partial\phi + c\partial\psi - c.c.)^n, \quad n \geq 0, \quad (7)$$

and have ghost dimension $2n$. An important feature[2,3,4] of $2d$ topological gravity is that (7) must be associated with a puncture operator

$$P = c\bar{c}\delta(\gamma)\delta(\bar{\gamma}) \quad (8)$$

of ghost dimension -2 . It creates two bosonic and two fermionic zero modes to give the right volume form on the moduli space of the punctured Riemann surfaces, so that there emerge non-vanishing correlation functions of local operators (7) times (8), representing the intersection numbers of the cycles on the moduli space[14,15].

The $D > 2$ topological string naturally appears as the topological gravity coupled with the above $c > 3$ topological matter, of which a covariant field-theoretical formalism can be obtained through direct extension of the $c < 3$ case[6]. Now the complete set of

nontrivial local physical operators, that are invariant under the total BRST operation of the coupled system, read

$$\sigma_{n,\alpha} = V_\alpha \cdot \sigma_n \cdot P \quad (9)$$

with V_α a covariantized version ($\varphi \rightarrow \varphi - c\lambda$, etc.) of (6), whose ghost charge (both left- and right-moving parts of the matter sector included) is $2n - 2 + 2\alpha/(k - 2)$. The correlation functions of operators (9)

$$\langle \sigma_{n_1, \alpha_1} \sigma_{n_2, \alpha_2} \cdots \sigma_{n_s, \alpha_s} \rangle_g \quad (10)$$

are defined as path integrals of them over the surface with g handles and s punctures and over the (supersymmetric) moduli space. The selection rule for non-vanishing values of (10) is just the ghost charge conservation. Taking into account the background charge of gravity sector associated with the supersymmetric moduli $-6(g - 1)$ and that of the matter sector from the $U(1)$ ghost current $2(g - 1)c/3$, it reads

$$\sum_{i=1}^s (n_i - 1 + \frac{\alpha_i}{k - 2}) = 2(g - 1) \frac{k - 3}{k - 2}. \quad (11)$$

In the meantime, the background charge of Liouville fields is cancelled by the BRST-invariant insertion of δ -function-surface-curvature creating operators $\prod_i e^{q_i} (\prod -c\partial\chi - \bar{c}\bar{\partial}\chi)$ with $\sum_i q_i = 2g - 2$ to the path integrals.

The amplitudes (10) are independent of the positions of operators, by the topological nature of the model. The integrals over these puncture positions (the moduli created by the punctures) do not contribute except in the region where punctures are very close or at least one puncture approaches a node on the surface (caused by the subtle compactification of the moduli space). From field-theoretical point of view, they constitute the so-called contact interactions. The analysis of them leads to a set of recursion relations for (10) in terms of genus g and the number of involved operators s . Let us simply present these relations with brief explanation. Detailed derivation is parallel to that in Ref.[4,6] where the techniques were developed.

First, the puncture operator P remains to behave as δ -function in the neighbourhood ϵ of another operator $\sigma_{n,\alpha}$: $\int_\epsilon P |\sigma_{n,\alpha}\rangle = |\sigma_{n-1,\alpha}\rangle$, such that

$$\int_\epsilon \sigma_{m+1,0} |\sigma_{n,\alpha}\rangle = |\sigma_{m+n,\alpha}\rangle. \quad (12)$$

Let the insertion of each curvature singularity $e^{q_{n,\alpha}(\prod -c\partial\chi - \bar{c}\bar{\partial}\chi)}$ with $q_{n,\alpha} = ((k-2)(n-1) + \alpha)/(k-3)$ be right at the position of $\sigma_{n,\alpha}$, the coefficient on the right side is then shifted to $\sigma_{n,\alpha} + 1$. Note by selection rule (11) alone, there could also exist a state $|\sigma_{m+n+1,\alpha-k+2}\rangle$ on the right side when $\alpha \geq k-2$. However, because the interaction from $\sigma_{m+1,0}$ is purely gravitational, no such state emerges. Eq.(12) gives rise to the contact terms $\langle \sigma_{m+1,0} \prod_i \sigma_{n_i,\alpha_i} \rangle_{\epsilon_j} \sim \langle \sigma_{m+n_j,\alpha_j} \prod_{i \neq j} \sigma_{n_i,\alpha_i} \rangle$, each of that represents the formation of a node separating two approaching points located on a sphere from the rest of the surface. The other type of contact interaction, occuring when a puncture is in the neighbourhood δ of a node, yields factorization terms $\langle \sigma_{m+1,0} \prod_{i \in S} \sigma_{n_i,\alpha_i} \rangle_{\delta,g} \sim \langle \sigma_{m_1,\alpha} \sigma_{m_2,\beta} \prod_{i \in S} \sigma_{n_i,\alpha_i} \rangle_{g-1}$ or $\langle \sigma_{m_1,\alpha} \prod_{i \in X} \sigma_{n_i,\alpha_i} \rangle_{g'} \langle \sigma_{m_2,\beta} \prod_{j \in Y} \sigma_{n_j,\alpha_j} \rangle_{g-g'}$ ($S = XUY$). Both indicate the formation of a second node, but the former breaks a handle whereas the latter divides the surface into two parts. Summing up these ingredients completes the integral over $\sigma_{m+1,0}$'s position, giving rise to the desired recursion relations:

$$\begin{aligned} \langle \sigma_{m+1,0} \prod_{i \in S} \sigma_{n_i,\alpha_i} \rangle_g &= \frac{k-2}{k-3} \sum_{i \in S} (n_i + \frac{\alpha_i - 1}{k-2}) \langle \sigma_{m+n_i,\alpha_i} \prod_{j \neq i} \sigma_{n_j,\alpha_j} \rangle_g \\ &+ \sum_{n=1}^m \sum_{\alpha=0}^k (b_{n,\alpha}^m \langle \sigma_{n-1,\alpha} \sigma_{m-n,k-\alpha} \prod_{i \in S} \sigma_{n_i,\alpha_i} \rangle_{g-1} \\ &+ \sum_{S=XUY,g'} d_{n,\alpha}^m \langle \sigma_{n-1,\alpha} \prod_{i \in X} \sigma_{n_i,\alpha_i} \rangle_{g'} \langle \sigma_{m-n,k-\alpha} \prod_{j \in Y} \sigma_{n_j,\alpha_j} \rangle_{g-g'}). \end{aligned} \quad (13)$$

From the selection rule (11), more factorization terms, such as $\langle \sigma_{n,\alpha} \sigma_{m-n,2-\alpha} \prod_i \sigma_{n_i,\alpha_i} \rangle_{g-1}$, are permitted; but they are not compatible with the contact terms, thus eliminated. Eq.(13) solves many correlation functions recursively upto the metric

$$\eta_{\alpha\beta} \equiv \langle \sigma_{0,0} \sigma_{0,\alpha} \sigma_{0,\beta} \rangle_0 = \delta_{\alpha+\beta,k} A_\alpha \quad (14)$$

and the one point function on the torus

$$\langle \sigma_{1,0} \rangle_1 = C. \quad (15)$$

The consistency of all equations in (13) and with (14)-(15), especially the requirement that positions of all operators in physical correlators are symmetric, determines all coefficients upto the normalization of $k+1$ primaries $\sigma_{0,\alpha}$:

$$b_{n,\alpha}^m = d_{n,\alpha}^m = \frac{k-2}{2(k-3)}, \quad A_\alpha = \frac{(\alpha-1)(k-\alpha-1)}{(k-2)(k-3)}, \quad C = \frac{(k^2-1)(k-6)}{24(k-2)(k-3)}. \quad (16)$$

Nevertheless, (13)-(16) are only partial solutions corresponding to two-point contact interactions initiated by pure gravity operator $\sigma_{m,0}$ and therefore involving spin-2 currents. The complete set of recursion relations correspond to multi-contact interactions of general $\sigma_{m,\alpha}$, which behave as higher-spin currents. Let us discuss them from the following constraint point of view.

Virasoro and W-Constraints — Consider the model's partition function

$$Z(t) = \langle e^{\sum_{n,\alpha} t_{n,\alpha} \sigma_{n,\alpha}} \rangle \equiv \sum_g \lambda^{2g-2} \langle e^{\sum_{n,\alpha} t_{n,\alpha} \sigma_{n,\alpha}} \rangle_g \quad (17)$$

perturbed by the couplings $t_{n,\alpha}$. It generates all unperturbed correlation functions

$$\langle \prod_{i=1}^s \sigma_{n_i, \alpha_i} \rangle = \frac{\partial^s}{\partial t_{n_1, \alpha_1} \partial t_{n_2, \alpha_2} \cdots \partial t_{n_s, \alpha_s}} \log Z(t). \quad (18)$$

The recursion relations (13)-(16) can be converted into the following Virasoro constraints

$$L_m Z(t) = 0, \quad m \geq -1 \quad (19)$$

with

$$\begin{aligned} L_{-1} &= \sum_{n=1}^{\infty} \sum_{\alpha=0}^k \left(n + \frac{\alpha-1}{k-2} \right) t_{n,\alpha} \frac{\partial}{\partial t_{n-1,\alpha}} + \frac{1}{2} \lambda^{-2} \sum_{\alpha=0}^k \frac{(\alpha-1)(k-\alpha-1)}{(k-2)^2} t_{0,\alpha} t_{0,k-\alpha}, \\ L_0 &= \sum_{n=0}^{\infty} \sum_{\alpha=0}^k \left(n + \frac{\alpha-1}{k-2} \right) t_{n,\alpha} \frac{\partial}{\partial t_{n,\alpha}} + \frac{(k^2-1)(k-6)}{24(k-2)^2}, \\ L_m &= \sum_{n=0}^{\infty} \sum_{\alpha=0}^k \left(n + \frac{\alpha-1}{k-2} \right) t_{n,\alpha} \frac{\partial}{\partial t_{n+m,\alpha}} + \frac{1}{2} \lambda^2 \sum_{n=1}^m \sum_{\alpha=0}^k \frac{\partial^2}{\partial t_{n-1,\alpha} \partial t_{m-n,k-\alpha}}, \end{aligned} \quad (20)$$

expanded around the critical point $t_{n,\alpha} = -\delta_{n,1} \delta_{\alpha,0}$. Note by these constraints the solutions can globally flow into other critical points corresponding to more general (topological) models.

Assuming (13)-(16) or (19)-(20) has a unique extension to the whole solution, the complete set of recursion relations are then equivalent to the (sufficiently large) extended W -constraints on $Z(t)$. To search for the latter, recall in the $D < 2$ case[6], the fact that the Virasoro constraints are the negative modes of the stress tensor realized with $k+1$ free bosons twisted by multiples of $1/(k+2)$ identifies the latter as that of level-1 $SL(k+2)$ current algebra. The relevant W -constraints then turn out to be the $SL(k+2)$

associated W_{k+2} algebra. For the present $D > 2$ case, (20) are nothing but the negative modes of the $c = k + 1$ stress tensor

$$T(z) = \sum_{\alpha=0}^k \tilde{j}_{\alpha}(z) \tilde{j}_{k-\alpha}(z) + \frac{(k^2 - 1)(k - 6)}{24(k - 2)^2 z^2} \quad (21)$$

in which $\tilde{j}_{\alpha} = \partial \tilde{\phi}_{\alpha} = \sum_n \tilde{j}_{n,\alpha} z^{-n-1-(\alpha-1)/(k-2)}$ are the free currents twisted by $(\alpha-1)/(k-2)$, with modes identified with $\tilde{j}_{n,\alpha} = \lambda \partial / \partial t_{n,\alpha}$, $\tilde{j}_{-(n+1),k-\alpha} = \lambda^{-1} (n + \frac{\alpha-1}{k-2}) t_{n,\alpha}$. Note it is straightforward to show that the weight vectors of the fundamental representation of graded Lie algebra $SL(p, q)$ have inner product

$$\vec{h}_i \cdot \vec{h}_j = \begin{cases} \delta_{ij} - 1/(p - q), & i, j = 1, \dots, p \\ 1/(p - q), & i = 1, \dots, p, j = p + 1, \dots, p + q \\ -\delta_{ij} - 1/(p - q), & i, j = p + 1, \dots, p + q, \end{cases} \quad (22)$$

matching the $1/(k-2)$ -twisting with $p = k, q = 2$. Thus the level-1 $SL(k, 2)$ current algebra appears to generate the relevant $c = k + 1$ W -algebra with (21) the Virasoro current, via standard reduction. The negative modes of this W -algebra's currents consistently serve as the desired W -constraints

$$W_{r,m} Z(t) = 0, \quad r \geq 2, m \geq -r + 1, \quad (23)$$

which are expected to exactly solve the $D > 2$ topological string.

$\hat{W}_{\infty}^{(p-q)}$ and $SL(p, q)$ — Let us further identify the W -algebra associated with $SL(k, 2)$. We proceed more generally. Recall it was shown[10] that there exists a nonlinear algebra $\hat{W}_{\infty}$, or even $\hat{W}_{1+\infty}$ by incorporating a spin-1 current, whose structure is most concisely exhibited as the second Hamiltonian structure of the integrable KP hierarchy with the spin- $(r+2)$ generators defined as the coefficient function $u_r(z)$ in the pseudo-differential Lax operator[16]:

$$L = D + \sum_{r=-1}^{\infty} u_r D^{-r-1}, \quad D \equiv \partial / \partial z. \quad (24)$$

Moreover, similar manipulation in terms of the powers of (24)

$$L^N = D^N + \sum_{r=-N}^{\infty} v_r D^{-r-1} \quad (25)$$

results in non-equivalent Hamiltonian structures $\hat{W}_{1+\infty}^{(N)}$ with $v_r(z)$ the spin- $(r + N + 1)$ generators[11]. The $\hat{W}_{1+\infty}^{(N)}$ ($\hat{W}_{1+\infty}^{(1)} \equiv \hat{W}_{1+\infty}$) algebra is proved[12] to admit an infinite number of free boson representations that $v_r(z)$ are realized through

$$L^N = \prod_{i=1}^p (D + j_i) \prod_{l=p+1}^{p+q} (D - j_l)^{-1}, \quad p - q = N, \quad (26)$$

as functions of free currents $j_i(z) = \partial\phi_i(z)$ and $j_l(z) = \partial\phi_l(z)$ of the opposite signature and their derivatives, generalizing the Miura transformation[17] for W_N and two boson realization[18] for $\hat{W}_\infty$. By imposing the (second-class) constraint $v_{-N} = \sum_i j_i + \sum_l j_l = 0$, $\hat{W}_{1+\infty}^{(N)}$ is reduced to $\hat{W}_\infty^{(N)}$. Correspondingly, (26) is modified to

$$L^N = \prod_{i=1}^p (D + \vec{h}_i \cdot \vec{j}) \prod_{l=p+1}^{p+q} (D - \vec{h}_l \cdot \vec{j})^{-1} \quad (27)$$

where $\vec{j} = (j_1, \dots, j_p, j_{p+1}, \dots, j_{p+q})$ and $\vec{h}_i$ are precisely the fundamental weight vectors of $SL(p, q)$ satisfying (22). Hence (27) is the Lax operator associated with $SL(p, q)$, via bosonic realization. Furthermore, the quantization of $\hat{W}_\infty^{(p-q)}$ may be carried out through, for instance, the Drinfeld-Sokolov reduction[19] from the level-1 graded $SL(p, q)$ current algebra. In particular, $\hat{W}_\infty^{(k-2)}$ fits the role of the relevant W -algebra from $SL(k, 2)$.

Remark — Finally, the exact solvability of the underlying $D > 2$ topological string suggests a close connection to complete integrable differential systems. Of prime consideration is the KP hierarchy. We speculate that the perturbed partition function (17) is a KP τ -function on its solution space[16], extending the wonderful outcomes at $D < 2$ [2,3,7,14,15] but within the Witten's general conjecture[2,14]. This τ -function is further expected to be written in terms of the Kontsevich-type matrix integrals[15], from which the fashion of $\hat{W}_\infty^{(k-2)}$ constraints would be more transparent.

Acknowledgements: The author thanks J. Lykken, M. Rocek, W. Siegel and Y.-S. Wu for discussions, and particularly C.N. Yang for providing a stimulating atmosphere and support. This work is supported in part by NSF grant PHY-9008452.

References

1. D. Gross and A. Migdal, Phys. Rev. Lett. 64 (1990) 127; M. Douglas and S. Shenker, Nucl. Phys. B335 (1990) 635; E. Brezin and V. Kazakov, Phys. Lett. B236 (1990) 144.

2. E. Witten, Nucl. Phys. B340 (1990) 281.
3. R. Dijkgraaf and E. Witten, Nucl. Phys. B342 (1990) 486.
4. E. Verlinde and H. Verlinde, Nucl. Phys. B348 (1991) 457.
5. E. Witten, Commun. Math. Phys. 117 (1988) 353; T. Eguchi and S.-K. Yang, Mod. Phys. Lett. A5 (1990) 1693.
6. K. Li, Nucl. Phys. B354 (1991) 711; 725.
7. R. Dijkgraaf, E. Verlinde and H. Verlinde, Nucl. Phys. B352 (1991) 59.
8. L. Dixon, M. Peskin and J. Lykken, Nucl. Phys. B325 (1989) 329.
9. W. Lerche, C. Vafa and N.P. Warner, Nucl. Phys. B324 (1989) 427.
10. F. Yu and Y.-S. Wu, Nucl. Phys. B373 (1992) 713; Utah preprint UU-HEP-92/12.
11. J. M. Figueroa-O'Farrill, J. Mas and E. Ramos, preprint BONN-HE-92/20; 92/ α .
12. F. Yu, to appear.
13. J. Lykken, Nucl. Phys. B313 (1989) 473.
14. E. Witten, Surveys In Diff. Geom. 1 (1991) 243.
15. M. Kontsevich, Funk. Anal. i Pril. 25 (1991) 50.
16. M. Sato, RIMS Kokyuroku 439 (1981) 30; E. Date, M. Jimbo, M. Kashiwara and T. Miwa, in Proc. of RIMS Symposium on Nonlinear Integrable Systems, eds. M. Jimbo and T. Miwa, (World Scientific, Singapore, 1983); G. Segal and G. Wilson, Publ. IHES 61 (1985) 1.
17. V. Drinfel'd and V. Sokolov, Sov. Probl. Math. 24 (1984) 81.
18. F. Yu and Y.-S. Wu, Phys. Rev. Lett. 68 (1992) 2996.
19. V. Drinfel'd and V. Sokolov, Journ. Sov. Math. 30 (1985) 1975.